EXERCISES: SESSION 1

- 1. Consider the following vectors of \mathbb{R}^2 : u = (1, 2), v = (1, 3), w = (-1, 2).
- (1) Compute a = u + v, b = u + 3v, c = 2v w
- (2) Represent u, v and w geometrically.
- (3) Represent a, b, c alongside their relationship with u, v and w geometrically.
- 2. Justify whether in each case $S \subset \mathbb{R}^3$ is a vector subspace:
- (1) $S = \{(0, 0, 0), (1, 0, 0)\}$
- (2) S is any finite collection of vectors of \mathbb{R}^3
- (3) $S = \{(x, y, z) \mid x + y = 0\}$
- (4) $S = \{(x, y, z) \mid x + y 1 = 0\}$
- (5) $S = \{v + w \mid v \in \text{span}\{(1, 0, 0), (0, 1, 0)\}, w \in \text{span}\{(1, 1, 1)\}\}$

3. Consider the following vectors of \mathbb{R}^2 : u = (1,3), v = (2,7), w = (1,5).

- (1) Are they linearly independent?
- (2) Do they span \mathbb{R}^2 ?
- (3) Compute two different linear combinations of u, v, w that produce the same vector b = (0, 1).
- (4) Are there more than one solution to the previous question?

4. Justify why the vectors (1, 0, 0), (0, 1, 1), (1, 0, 1) are a basis of \mathbb{R}^3 .